



PAXLOVID®

Patient Information

This information leaflet includes important information about the medicine called **Paxlovid®** when used in the treatment of COVID-19. Having COVID-19 infection and being treated does not offer the same protection as vaccination.

What is the medication used for?

Paxlovid® has been provisionally registered by the Therapeutic Goods Administration in Australia to treat adults with COVID-19 who do not need oxygen but are at risk of severe illness and hospitalisation. This medicine is not intended to be a replacement for vaccination against COVID-19.

Paxlovid® is used in adults to treat mild to moderate coronavirus disease (COVID-19) in patients:

- who have a positive result from a severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) viral test, AND
- who have a high risk of getting severe COVID-19, including hospitalisation or death.

Paxlovid® must not be used:

- to treat patients with severe or critical COVID-19 infection
- to treat patients who have had symptoms for more than 5 days
- as a preventive medicine to stop COVID-19 infection from occurring
- for longer than 5 days in a row
- to treat children and adolescents less than 18 years of age.

Do not give your medicine to anyone else, even if they have the same condition as you.

What does this medication do?

COVID-19 is caused by a virus called a coronavirus. Paxlovid® contains 2 antiviral medicines that are co-packaged together, nirmatrelvir and ritonavir.


Paxlovid® stops the virus from multiplying. This can help your body to overcome the virus infection and may help you to get better faster.

Paxlovid® consists of 2 medicines co-packaged together:

- **Nirmatrelvir (pink tablet):** 150 mg
- **Ritonavir (white tablet):** 100 mg

What should be considered when using Paxlovid® for COVID-19?

Paxlovid® is a new medicine so information about how well it works and how safe it is, is being collected



It is important that patients and their carers understand when and why Paxlovid® may be useful. Your health practitioner will provide more information about how well it works and how safe it is for your condition.

The use of Paxlovid® for COVID-19 is new and it is important for you to provide your formal consent before Paxlovid® treatment begins. You can always change your mind about treatment with Paxlovid® and withdraw your consent at any time.

How do you store the medication?



Store at room temperature (15°C to 30°C) away from sunlight, excessive heat and moisture. Keep out of reach and sight of children and pets.

How should this medication be used?

- Paxlovid® consists of 2 medications: nirmatrelvir and ritonavir.
- Take 2 pink tablets of nirmatrelvir with 1 white tablet of ritonavir by mouth twice a day, in the morning and at bedtime, for 5 days. Take all 3 tablets at the same time
- You must always take the nirmatrelvir tablets at the same time as the ritonavir tablet.
- If you have kidney problems, you may need to take a lower dose. Check with your health practitioner.
- You can take Paxlovid® with or without food. Swallow the tablets whole. Do not break, chew or crush the tablets.
- Even if you feel better, it is important to take the full 5-day treatment course. Do not stop taking Paxlovid® without talking to your health practitioner first.
- Talk to your health practitioner if you do not feel better or if you feel worse after 5 days.

Pharmacist or health practitioner to tick relevant regimen.

☐ **STANDARD DOSE** – Take 2 pink tablets of nirmatrelvir with 1 white tablet of ritonavir by mouth every 12 hours at the same time each morning and evening. Use this calendar to help keep track of your doses. Note, if your first dose was in the evening on Day 1 your last dose may be in the morning on Day 6.



	Day 1	Day 2	Day 3	Day 4	Day 5
	Date:	Date:	Date:	Date:	Date:
Morning  2 x nirmatrelvir 1 x ritonavir Time:					
Evening  2 x nirmatrelvir 1 x ritonavir Time:					

What should the health practitioner know before Paxlovid® is used in COVID-19?

The health practitioner should know about:

- any medical conditions you have
- previous allergic reactions to any medicine
- all medicines including over-the-counter and complementary medicines e.g. vitamins, minerals, herbal or naturopathic medicines that you are taking or have recently taken
- the possibility of pregnancy or plans for pregnancy
- recent vaccinations or plans to get vaccinated

☐ **REDUCED DOSE (if you have reduced kidney function)** – Take one (1) pink tablet of nirmatrelvir with one (1) white tablet of ritonavir by mouth every 12 hours at the same time each morning and evening. **The pharmacy will modify the packaging of the medication to ensure you have the correct number of tablets for your dose.** Use this calendar to help keep track of your doses. Note, if your first dose was in the evening on Day 1 your last dose may be in the morning on Day 6.

	Day 1 Date:	Day 2 Date:	Day 3 Date:	Day 4 Date:	Day 5 Date:
Morning  1 x nirmatrelvir 1 x ritonavir Time:					
Evening  1 x nirmatrelvir 1 x ritonavir Time:					

What should you do if you forget a dose?

- If it has been less than 8 hours since the missed dose, take the dose as soon as you remember.
- If it has been more than 8 hours since the missed dose, skip this dose and take your dose at the next scheduled time.
- Do not double the dose to make up for a missed dose.

What should you do if you take too much?

If you think you, or a person you are caring for, has taken too much Paxlovid®, you may need urgent medical attention.

You should immediately:

- contact your health practitioner OR
- phone the Poisons Information Centre (13 11 26), OR
- go to the emergency department of your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

What are some common side effects?

Paxlovid® is generally well tolerated – most people only experience mild side effects.

Rarely, Paxlovid® may cause liver problems, or resistance to HIV medications if you have untreated HIV infection.

Some other reported side effects include:

- altered sense of taste or metallic taste in mouth
- vomiting or diarrhea
- high blood pressure
- muscle aches
- headaches

Unwanted side effects are possible with all medications. The effects listed are not all of the side effects that can occur.

If you have a troublesome symptom or side effect that is not listed here or becomes severe enough to interfere with your daily activities, tell your health practitioner.

Speak with your healthcare provider if you have questions about side effects.

If you are experiencing any side effects, including any of the ones listed above or are feeling unwell in any way, please call the health practitioner on the phone number below for advice.

Name of health practitioner to call:

Telephone number:

Who should I know before I take Paxlovid®?

- **Do not take Paxlovid® if you are allergic to nirmatrelvir, ritonavir or to any of the other ingredients in Paxlovid®:**
 - Nirmatrelvir tablets: microcrystalline cellulose, lactose monohydrate, croscarmellose sodium, colloidal silicon dioxide, sodium stearyl fumarate.
 - Ritonavir tablets: copovidone, calcium hydrogen phosphate, sorbitan monolaurate, hypromellose, colloidal anhydrous silica, titanium dioxide, sodium stearyl fumarate, macrogol 400, hypromellose, purified talc, macrogol 3350, polysorbate 80.
- **Tell your health practitioner if you are pregnant, think you might be pregnant or are planning to become pregnant.** You should not take Paxlovid® if you are pregnant unless your health practitioner advises that you can. Women should use effective contraception while they are taking Paxlovid®. Paxlovid® may affect how birth control pills, patches and vaginal rings work. You should use alternative contraception or an additional barrier method such as a condom while you are taking Paxlovid®. Talk to your health practitioner about effective methods of birth control.
- **Breastfeeding:** Tell your health practitioner if you are breastfeeding or plan to breastfeed. Paxlovid® can pass into your breastmilk. Your health practitioner will tell you if you can breastfeed your baby while taking Paxlovid®.
For further information about the use of this medicine in pregnancy, breastfeeding or family planning, please contact the King Edward Memorial Hospital Obstetric Medicines Information Service on 6458 2723.
- **Patients with kidney problems:** Tell your health practitioner before you take Paxlovid® if you have any kidney problems. You might need a lower dose of Paxlovid®. Your health practitioner will prescribe a dose that is right for you.
- **Liver problems:** Before you take Paxlovid® tell your health practitioner if you have any liver problems. Liver problems have happened in patients taking ritonavir, a medicine in Paxlovid®. Talk to your health practitioner if you get any symptoms of liver problems. These

may include yellow skin or whites of eyes, nausea, tiredness or feeling unwell, loss of appetite, fever, skin rash, abdominal pain, pale stool or dark coloured urine.

- **Paxlovid® interacts with many medications and herbal supplements.** Taking Paxlovid® with some medicines may cause serious or life-threatening side effects.

Let your doctor know if you are taking, or have recently taken, any of the following medicines as they should not be taken with Paxlovid®:

- **alfuzosin** - medicine to treat an enlarged prostate
- **pethidine, piroxicam** - medicines to relieve pain
- **amiodarone, flecainide** - medicines to correct or change heart rhythm or lower blood pressure
- **neratinib, apalutamide, venetoclax** - medicines to treat certain types of cancers
- **colchicine** - medicine to treat gout
- **lurasidone, clozapine** - medicines to treat certain mental and emotional health conditions
- **ergometrine** - medicine to stop excessive bleeding that may occur following childbirth, miscarriage or termination of pregnancy
- **simvastatin** - medicine to lower blood cholesterol
- **avanafil, vardenafil** - medicines for erectile dysfunction
- **sildenafil** - medicine for erectile dysfunction, or to treat high blood pressure in the lungs
- **tadalafil** - medicine for erectile dysfunction, or urinary difficulties due to an enlarged prostate
- **diazepam** - medicine to assist anxiety, agitation or muscle spasms
- **carbamazepine, phenobarbital, phenytoin** - medicines to treat epilepsy or prevent convulsions.
- **rifampicin** - medicine to treat tuberculosis
- **St. John's Wort** (*hypericum perforatum*) - a herbal remedy used for depression and anxiety.

- This is not a complete list of medicines that interact with Paxlovid®.
- Tell your healthcare professional about ALL of the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines
- Your doctor will check they are suitable to take with Paxlovid®, as some medicines may need extra monitoring, a dose change or be stopped while you are taking Paxlovid®.
- Talk to your healthcare professional first before taking any new medicines while taking Paxlovid®. They will tell you if it is safe to take.

What should I do if my COVID-19 symptoms get worse?

Call your healthcare provider if you have any questions or concerns.

Call 000 immediately if you experience severe symptoms of COVID-19, such as:

- difficulty breathing
- severe chest pain
- feelings of disorientation or confusion
- loss of consciousness.

If you have any questions about this medication, call your health practitioner.

What should I do after I finish my treatment?

Treatment does not shorten your isolation time. You are still considered to be contagious while you are taking your treatment. Follow all Public Health guidelines for isolation.

For information about isolation please refer to the Healthy WA website.

https://www.healthywa.wa.gov.au/Articles/A_E/Coronavirus/Quarantine-and-isolation

This document can be made available in alternative formats on request for a person with disability.

© Department of Health 2022

Copyright to this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the *Copyright Act 1968*, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.

Adapted from The Government of Canada, *Drug Details: Paxlovid* <https://covid-vaccine.canada.ca/info/paxlovid-en.html>